



This page from the Epilepsy Foundation website can be viewed at:

<http://www.epilepsyfoundation.org/epilepsyusa/news/Study-Confirms-Favorable-Long-term-Prognosis-of-Epilepsy.cfm>

Study Confirms Favorable Long-term Prognosis of Epilepsy

A study conducted by researchers in The Netherlands confirmed that children with idiopathic (of unknown cause) new-onset epilepsy have a significantly higher rate of remission than those with remote symptomatic epilepsy. Results of this study are now available online in *Epilepsia*.

“Knowledge of the long-term outcome of childhood epilepsy is an important factor in treatment decisions,” explained study leader Ada Geerts. “The course of epilepsy is in our view important to investigate, because intra- and inter-individual variation may be considerable. Periods of remission and relapse may interchange, which may influence our interpretation of the results of prognostic studies, advice to patients, treatment strategies and timing of referral for surgery.”

Researchers indicate it is still largely unknown whether the long-term outcome of epilepsy is determined by its natural course or whether it can be modified by treatment. Whereas outcome in terms of remission or death can be determined easily, the long-range course of epilepsy is more difficult to describe. To address the need for studies with prolonged follow-up in a large group of patients, the research team evaluated the course and outcome of childhood-onset epilepsy in 413 children over a 15-year period.

The study began in 1988 with the team evaluating 494 children (aged 1 month to 16 years) who were newly diagnosed with epilepsy. The patients were followed for 5 years, and then contacted again by questionnaire 10 years later. A total of 413 responses were received.

Among the people who responded, 70.9 percent had a terminal remission of at least 5 years at the end of the study. In contrast, epilepsy remained active in 30 percent of patients and became intractable (non-responsive to treatment) in 10 percent, with the majority of those with active epilepsy or intractability having a non-idiopathic etiology. In patients with a favorable course, idiopathic etiology was predominant, and the mean longest remission during the first 5 years of follow-up was significantly higher than in all other groups.

The researchers concluded that antiepileptic drugs (AEDs) probably do not influence epilepsy’s course—they merely suppress seizures. Mortality was significantly higher only in those with remote symptomatic etiology.

Geerts concluded, “The long-term prognosis of epilepsy is favorable in the majority of children, especially for those with idiopathic etiology. It remains to be seen whether such a course is influenced by the treatment given, since childhood-onset epilepsy is often a benign, self-limiting disorder and treatment proved to be ineffective in those with active epilepsy or intractability.”